Red 1/20/00 upl

ACCEPT 70

COLORADO RARE BIRD REPORT

MOURNING WARBLER
Oporornis philadelphia

Location: Lamar Community College grove = "Willow Creek"; Prowers County; Lamar, Colorado

Date:

May 9-15, 1999

Setting:

We first found this bird with remnants of the May 8th CFO field trip. The bird was well seen on May 9th and then subsequently seen for almost a week. Although it started as a very secretive warbler, there were many occasions when the bird foraged quite openly and gave "crippling" good views. On the 9th I observed it at first from 8:00-8:30 a.m. from distances of 25-40 feet with B&L Elite 10x40 binoculars. The day was sunny. It foraged most often from ground level to up to mid-levels of 10 feet or so. It moved stealthily at times, like Oporomis sp. warblers. The bird gave a flat, dry "chip" note rarely, but especially when it flushed

Description:

The bird was a breeding-plumaged male. It was about the size, shape and pattern of a MacGillivray's Warbler, being a large warbler — about 6" long and was quite heavy-bodied and long-tailed. It was uniformly olive colored above. The wings were dark and lacked wingbars. It had a complete dark, gray hood that came down across the chest. The hood was black at the bottom. The undersides were otherwise a bright, deep yellow, including the undertail coverts. The face was dark gray, like the crown and hood — and the blackest part of the face was a blackish smudge around the eye. There was no trace of white eye crescents nor eyering.

Similar Species:

The only other true ID contenders are MacGillivray's Warbler (smaller bib; dull to bright white eye crescents) and Connecticut Warbler (less extensive, lighter gray hood; complete white or buff eyering; gray, not black, lores).

Experience with Species:

I have seen all Oporomis sp. in Colorado and/or elsewhere in the U.S. The most recent Mourning Warblers that I have observed were 9 during 1979 in northern Illinois and a in September 1993 at Lake Henry and a male in Colorado in 1995.

This report is prepared from notes made during and after observation.

Mark Janos

10 Sedum Ct. Pueblo, CO 81001 Marlyanes

CFO-RC FILE # 1999-150

ACCEPT 7-0

Red 5/99 upl

COLORADO FIELD ORNITHOLOGISTS SIGHT RECORD FORM

RESULTS
PLEASE PRINT OR TYPE. ATTACH PHOTOGRAPHS AND ADDITIONAL PAGES IF NECESSARY.
SPECIES Morrang Warbler Operonis philadelphia (SCIENTIFIC NAME)
DATE(S) OF OCCURRENCE MAY 8 1996
NUMBER OF BIRDS SEENOne_ SEX_mAle_PLUMAGE_AlterNATE_
EXACT LOCATION South end of Riporians glove on Willow Creek AT LAMAR Com. College
COUNTY Prawers NEAREST TOWN LAMAR ELEVATION 3622
LENGTH OF OBSERVATION 3 min TIME OF OBSERVATION - 9:00 Am
OTHER OBSERVERS: MARK JANUS, BRANDEN K. PERCUAI, DAVE QUESENBERRY VIL ZERBIE BUB DICKSON, MO. B.
LIGHT CONDITIONS - SURMY, CLEAR SK.ES
OPTICAL EQUIPMENT USED: ZIES> 10 x 40
DISTANCE FROM THE BIRD: 10 FEET
WAS THE BIRD PHOTOGRAPHED YN PHOTOGRAPHS SUBMITTED Y/N WHO TOOK THE PHOTOGRAPHS?
DESCRIBE THE GENERAL AND SPECIFIC HABITAT IN WHICH THE BIRD OCCURRED: Willow CREEK RIPARIAN AREA in dense willows Along
the CREEK AND in A willow, golden correct thicket NEAR the MAINTAINE GARAGE.

DESCRIBE THE BIRD'S BEHAVIOR (FLYING; FEEDING; HABITS; BEHAVIOR USED IN IDENTIFYING THE SPECIES OR IN SUPPORTING THE ID):

BIRD WAS 15 OBSERVED SUKING AT the BASE of Willows Along Willow CREEK.

Hopping on ground And Amongst the base of the Thickess. LATER THE BIRD WAS

SEEN WIRING At EYE LEVEL in A Willow, golden CURRENT Thicket

DESCRIBE THE BIRD IN DETAIL. THIS IS THE MOST IMPORTANT PART OF THIS REPORT. INCLUDE OVERALL SIZE AND SHAPE, AND HOW YOU PLACES THE BIRD IN THE GENERAL GROUP THAT YOU DID. GIVE COMPLETE DETAILS OF PLUMAGE, INCLUDING HEAD, THROAT, UNDERSIDES NAPE, BACK WINGS, AND TAIL. GIVE PARTICULAR EMPHASIS TO MARKS YOU USED TO ID. THE BIRD. DESCRIBE OTHER DETAILS SUCH AS COLOR AND SHAPE OF BILL, LEGS, EYES, ETC. THAT WERE IMPORTANT TO MAKING THE ID.. (ATTACH ADDITIONAL SHEETS, DRAWINGS, SKETCHES, OR COPY OF FIELD NOTES):

I first observed AN operanis worder suking in the under brush. Checking to see if it was a MacGillivary's worder I noticed the bird HAD NO White eye CRESTS. Upon Father observation I SAW A

SLATE gamy head with Brank Eyes who Eye CRESTS. The slate

9RAY of the NAPE -changed to olive green on the BALK & Rump & TAIL

\$ Wings. Wings were with out wing bars. The gray of the

Wings. Wings were with out wing bars. The gray of the

head extended on to the breast to form a BiB. Black feathers

head extended on to the breast to form a BiB. Black feathers

Filled the bib from the the chin through the threat and wides

At the base of the bib. The Berly & under toil courses & blanks

WERE A Lemon YEllow was stellight a grayish how on the flanks

MACG. 11. VRAY'S WARDER - to White EYE CRESTS BLACK IN BIB NOT AS EXTENSIVE
COMETICUT WARBIER - White EYE Ring LACKS BLACK IN BIS WATHER THAN hops
WHAT IS YOUR PRIOR EXPERIENCE WITH THIS AND SIMILARLY APPEARING SPECIES:
LIST BOOKS, GUIDES RECORDINGS, OR OTHER SOURCES CONSULTED AND HOW THIS INFLUENCED YOUR ID. (DURING AND AFTER OBSERVATION):
THIS REPORT WAS MADE FROM NOTES MADE DURINGOR AFTER OBSERVATION, OR LATER FROM MEMORY_V
DATE OF WRITTEN REPORT 5-18-99
REPORTING OBSERVER: RIC OLSON
SIGNATURE ROL
9450 N. HEATHER DR. CASTLE ROCK, CO. 80104

303-790-7753

OLY@ECENTRAL.COM