Species: PIPING PLOVER	ORD Accepted GR "/ Charadrius melodus	
(Vernacular Name)		
	(Scientific Name-use b	
Date(s): September 29, 1974 Time B	ird Seen:to	
Locality: Jackson Reservoir		
Nearest Town: Goodrich	County: Morgan	
Other observers who identified this bird:		
Jack Reddall 4450 S. Alt	DRESS	TH
4470 S. AT	on St., Englewood	77
(missel Posterot C V colder		
Optical Equipment: 7 X 50 binoculars; Bausch Light Conditions: Sunny, with sun coming from	and Lomb 15X-60X Zoom Telescope	
Distance from Bird (how measured):		
Number of Birds Seen: One Sex: Unknown Overall Appearance and Size: A small plover,	Plumage: nonbreeding	
- brown upperparts, incomplete breast ring an	d orangish legs.	.s. p
	The state of the s	
Detailed Description (describe only what was	actually observed in the field):	
Bill (shape and color): short and stubby, bl Legs (shape and color): average in length, y	allow or vallow orence in color	
Crown and Forehead: forhead whitish, crown pale	brownish.dark mark on fh me (7
hape:_pale brown Face: pal	e brownish-white Throat. Whi	te
breast and Sides: white, with incomplete pale	brownish breast ring on the side	s of
Belly and Flanks: white Undertail Coverts: white		
Back: pale brown		
Wings: pale brown with whitish wing stripe		
Underwings: not seen		-
Rump: pale brown Upper Side of Tail: brown, becoming darker to	Uppertail Coverts: whitis	h
Under Side of Tail. not seen		
Shape of Tail: average in width, fairly short	, rounded	
Voice: no notes heard		
A A Parada Barata proper		
Behavior: observed walking and feeding in mud	flats at edge of the water, but	not
into the water. Also seen preening and spreading that at edge of large plains	eading wing, but not seen in flig	ht
Prior Emerience with this Species: Five obse	rved on August 24. 1970 et Wilder	ness
Park in Emmet County, Michigan.	.//0 60 111061	
Total Signa Circles Assessment Control	od? Sandnings + 11-1-1-1	
TOW WELL DIMINAL A DUNDONS NO SUCCESS ELS	untain Ployer s eliminated due to	amo
How Were Similarly Appearing Species Eliminate and bill. Killdeer, Black-bellied Golden Mon	hill instead of this 1222	re in
Snowy eliminated because of short and stubby	THE THE TERM OF THE PARTY AND THE	
Snowy eliminated because of short and stubby black legs, whitish instead of brownish rum	. dark hand at end of the toil on	d_{-th}
Showy eliminated because of short and stubby black legs, whitish instead of brownish rump likelihood of a Piping at that locality at the	, dark band at end of the tail an	+
Snowy eliminated because of short and stubby black legs, whitish instead of brownish rum	, dark band at end of the tail an	+
Snowy eliminated because of short and stubby black legs, whitish instead of brownish rump likelihood of a Piping at that locality at the of strong breast band which Semipalmated usus upperparts.	, dark band at end of the tail an hat season. Semipalmated elimina ally has in fall, and much lighte	ted r br
Showy eliminated because of short and stubby black legs, whitish instead of brownish rump likelihood of a Piping at that locality at the of strong breast band which Semipalmated usus upperparts. This Report was Written From: Notes made during the strong breast band which semipalmated usus upperparts.	, dark band at end of the tail an hat season. Semipalmated elimina ally has in fall, and much lighte	ted r br
Snowy eliminated because of short and stubby black legs, whitish instead of brownish rump likelihood of a Piping at that locality at the of strong breast band which Semipalmated usus upperparts. This Report was Written From: Notes made during the strong breast band which semipalmated usus upperparts.	, dark band at end of the tail an hat season. Semipalmated elimina ally has in fall, and much lighte	ted r br

COLORADO FIELD OPUTTHOLOGISTS SIGHT RECORD REPORT [X] RARE OR UNUSUAL RECORD [] HEW STATE RECORD.
Species: PIPING PLOVER
Species: PIPING PLOVER Charadrius melodus (Vernacular Name) (Scientific Name-use binomial)
Date(s): September 29, 1974 Time Bird Scen: 1:45 P.M. to 2:15 P.M MDT
Locality: Jackson Reservoir
Nearest Town: Goodrich County: Morgan Cther observers who identified this bird:
NAME ADDRESS Robert Andrews - 1390 South Ivy Way, Denver, Colorado 80222 TELEPHONE
Cotical Equipment: 10X50 Bushnell Binoculars; 15X60X Zoom Telescope Light Conditions: Bright, clear; Sunlight from behind observers; air still; slight heat Distance from Bird (how measured): 100 feet to 250 feet, estimated. /// waves Murber of Birds Seen: One Sex: Unknown Plumage: Fall (non-breeding) Overall Appearance and Size: Small, extremely pale plover-like bird approximately the same size as many nearby Baird's Sandpipers. Much smaller than many nearby Killdeers and much paler. Appeared more chunkier than the many "peeps" with which it was feeding. Detailed Description (describe only what was actually observed in the field): Bill (chape and color): Small, stubby; dark appearing. Legs (shape and color): Noticeably yellowish, short legs. Feet not noted. Croum and Forchead: Forehead white; Croum very pale tan Eye (color): Dark Mape: Very pale tan Face: White, slight ear patch Throat: White Breast and Sides: White with incomplete, single neck band of pale tan in color. Belly and Flanks: White Undertail Coverts: White Back: Pale tan Wings: Tan with dark primaries Underwings: Not observed as bird did not fly.
Pump: White, unmarked Uppertail Coverts: Not noted Upper Side of Tail: White, marked with dark brown on lower half Under Side of Tail: Not seen
Shape of Tail: Short, rounded, un-notched
Voice: Silent
Behavior: Typical plover actions - running along mud flat in short spurts, pausing now and then, occasionally picking up bits of food. Habitat: Mud flat along shore of fresh water reservoir on high plains. Prior Experience with this Species: Have seen twice before in Colorado, both fall records from Jackson Reservoir. Over twenty years experience from Long Island, New York area
where it breeds regularly and commonly on the outer beaches.
How Were Similarly Appearing Species Eliminated? Snowy Plover has longer, thinner bill and black or dark legs. Snowy Plover has a rather conspicuous dark ear-patch not evident in the Piping Plover. Although the breast band on the fall Piping Plover that we saw was incomplete, it was far more prominent and reached considerably further down the sides than is evident in the Snowy Plover. Semipalmated Plover is much darker, even in the fall, on the upper parts and has prominent white forehead and eyeline. Wilson's Plover is much larger, darker with booms bill (and best prominent white forehead and eyeline. Wilson's Plover is much
larger, darker with heavy bill (and has never been reported in the state). Killdeer is much larger still, darker and has two breast bands. This Report was Written From: Notes made during X afterobservation; Memory
Date Prepared: October 2, 1974 Town and State Englewood, Colorado ZTP 80110
Return to. TACK DUDDATT 11 do c 11 121