COLORADO FIELD ORNITHOLOGISTS - SIGHT RECORD REPORT	
Species: SCOTTUS_ORIOLE (Vernacular Name)	(Scientific Name-use binomial)
Date(s): May 3 1075 Time Bird	Seen: _9:05 to _9:30
Nearest Town: Littleton	County: Arapahoe
Other observers who independently identified this ADDRESS	TELEPHONE
ng Kingery, Thompson Marsh, Ralph Sargent, Doug & Ju Wilbur, Ted Haase, Joe Rolder, Tom Kerwin, Robert	
Floyd Pendell Optical Equipment: <u>7 x 35 binocs; Thompson H</u> If photographed, type of equipment used:	
Light Conditions: <u>sunny, good light</u> Distance from bird (how measured): 50 feet	
Number of Birds Seen: 1 Sex: Plum	age:adult

Describe the bird's overall behavior and demeanor:

U

The bird flew into the top of a cottonwood tree and fed, at the top, for the Balf hour our necks could watch it. It picked insects (?) off the catkins, pirouetting around them and showing all of its field marks--except it managed to keep the top of the tail concealed from view most of the time.

Describe in great detail the bird's size, overall shape, plumage and color-pattern. Include details on shape of <u>bill</u>, wings and <u>tail</u>. Give paticular emphasis to the marks you used to identify the bird. Reference to its resemblance in field guide descriptions is not enough. Describe what you actually saw in the field:

Size of a robin; oriole shape, with the typical long pointed oriole bill. It had a black head--hood actually, as it came down to cover the upper chest, a **back** black back, black wings with one white wing bar and a yellow wedge at the base of the wings--formed by the black wings, black back, and black hood. The tail was black on top, with a narrow yellow edges on the upper half. The bottom of the tail was black on the ventral side of the tail half black, **the the dot** (see drawings). The yellow base on the ventral side of the tail lapped over the edge of the dorsal side of the tail, providing the narrow yellow **back** nd on the upper half of the dorsal side. The lower chest and breast and belly were also yellow. The yellow was a bright lemon-typed yellow color. The hood was ragged-across the breast--i.e., it did not have a definite, even line between the two colors.

And see descriptions of other observers, attached.

\* the yellow on the top of the tail was not nearly as conspicuous as Robbins shows; it was closer to Peterson in Mexican birds, or perhaps even less noticeable.

Slar le (OVER)

Describe the bird's song or call, if heard, including method of delivery, i.e., from perch, in flight, etc:

No sound.

## Describe the habitat in which you observed this bird:

Cottominwood river bottom, in a cottonwood with catkins in full bloom (or whatever they do). One hundred yards below the Waterton bridge, beside the South Platte River.

# How were similarly appearing species eliminated? Bright lemon yellow and the cripp black hood

The white wing bar, black back, and yellow on the tail eliminate all the Mexican species, including the Black-headed.

# Prior experience with this and similarly appearing species:

I've seen the Scotts in Arizona, California, and Nevada several times, including about a month ago. I haven't seen any of the Mexican ones, and the second s

Books, illustrations and advice consulted and how did these influence this documentation:

We all used Birds of North America, and learmed what field marks to look for to eliminate similar species. A review of Peterson's Mexican Guide helps in checking the other Mexican species.

This report was written from notes made during XX after XX observation; from memory

PRINT NAME: Hugh Kingery	SIGNATURE A A A DATE.	5/9/75
Street Address:869 Milwaukee	SIGNATURES DATE: Town and State Penver Colo 80	206 ZTP:
Return to: JACK REDDALL - 4450 South AD		

Field notes by observers of Scott's Oriole, May 3, 1975.

54-75-90 gr 9/26/25

## Urling Kingery

Black head attached to black back. Yellow wing patch like bar, white wing bar below that. Yellow breast. Yellow bars down sides of tail.

#### Joe Roller.

Size and shape of Northern Oriole. Seen 15 min., 50 feet in top of tree. Great light, 7x50 glasses. "Goldfinch" yellow sides and belly. Black head, eye, back and wings. Rump yellow. White wing mark. Tail black above, yellow below, black tip below. Black of back extends down back farther than on picture in Robbins, making rump look smaller.

## Andy Wilbur

Yellow underside, black head, yellow comes down on back of tail, yellow comes underneath and down on tail. So ight wing bar.

#### Judy Ward

Black head. Bright yellow body (yellow like Am. Goldfinch). Yellow up toward rump. Undertail coverts yellow. Undertail black tip. One white wing bar. Yellow "bar" in front shoulder. Dark back.

Doug Ward (Age 9)

Black head, black back and wings and Redstart tail pattern. White and yellow wing bars. Time seen, 9:15.

### KanxKernin

R. Spencer '

Cottownwoods, feeding catkins.

Black bill, black hood, yeldow belly, yellow halfway down undertail. Large yellow wing bar. White wing bar. Black wings & tail. 9:05--9:20 a.m.

(C	54-75-90
COLORADO FIELD ORNITHOLOGISTS SIGHT RECORD REPORT RARE OR UNUSUAL RECORD . NEW STATE RECORD	CFO-ORC FILE # Accepted gR 9/26/75
Species: SCOTT'S ORIOLE	Icterus pavisoviem
(Vernacular Name)	(Scientific Name-use binomial)
Pote(s): MAY 3 1975 Time Bird Seen	: 9:10 Am to 9:25 AM
Lucality: WATERTON COLORADO . 50.	-75 VIS HOWNSTREAM -BRIDGE-D FLANE
Nearest Town: WATERTON	County: JETFERSON
Other observers who identified this bird: NAME ADDRESS	TTOUCHE
THOMPSON MARSH -	
others uptical Equipment: ZETSS 10x40	ZIAD HICHTOPOFONTONIZE
Distance from Bird (how measured): 100 FT C	
Number of Birds Seen: <u> </u> Sex: <u>M</u> Overall Appearance and Size: <u>ORIGLE TYPE</u>	Plumage: ADULT SPRING
Detailed Description (describe only what was actual Bill (shape and color):ONG STRAIGHT - DAR Legs (shape and color):ONGT OFFICTUE Crown and Forehead:OLACK_ Nape:FLACK Breast and Sides: Belly and Flanks:	Eye (color)_DARK
Undertail Coverts: VELOW	
Back: BLACK - VELLOW ON SHOULDER	SLICHT WHITE PATCHES
Underwings: DANAT OBSCRUE	
Bumn: VITIAN	Uppertail Coverts: VHT Low
Upper Side of Tail: JELLOW . Farek -VET	JON ON ERIAN FERMINERS
Under Side of Tail: YELLOW + PLACE - Y. Shape of Tail: Long - Wide	<u>ELDIC ONE APPE ABCONTO CONF</u>
Voice: DIDNOF HEAR	
Charton: BIRD MONIALE APPRIAID AT TO	OP OF MATURE COTTONIOGOD - SOMETIME
QUITE TTILL SO OPSERVATION VERY CLE	AR.
Habitat: AT TOP OF MATURE COTTONILOOD A	LONG SOUTH PLATTE
Prior Experience with this Species: ALANE	
Wow Were Similarly Appearing Species Eliminated? RIOLE, BLACK-HEADED HAS VELLOW BA	CK. SCOTTSORIOLE FLACK BACK - 11115
TAIL - BLACK SCOTT'S YELLOW OUTER TAIL DIFFERILE CLEARLY OBSERVED ON UN	FEATHERS HAIFLENGHT OF TAIL 1415
OBSERVED ON UPPERSIDE OFTAIL BE	N ACROSS WHOLE TAIL. LESS DISTINCTLY
This Report was Written From: Notes made during-	- after X observation: Memory X
	Part Part Com
Ralph Sargent	

				54-75-90	
	ITHOLOGISTS - SIGHT RE		CTO CRO FTLE	10. XX	_
RARE OR UNUSUA	l record []nem s	STATE RECORD	ACCEPTED	REJECTED /R 9/2	17
Secies: - Secti	Minto 3		Icterus pari	DATUM	
Species: - Maria	(Vernacular Name)			Name-use binomial	)
Date(s): 5-3-75 Iccality: Water to	A we were and	and the second of the		to come 9:20	
Nearest Town:			County:	forson	-
	no independently ident:	ADDRESS	• • • • • • • • • • • • • • • • • • •	TELE	PH
Boo Apen	ier				_
	7 X 50 and 20 X 60				
Light Conditions:	good	and the state of the			
Distance from bir	d (how measured): 40-	50 feet estimate	a (top y locg co	ttonwood)	
					-
Number of Birds S	een:l_Sex:	7 Plumage:	aduct		
	's overall behavior an		ld moved a	bout occasionally	
Man the tra	of a beg cottonwood	a			

Describe in great detail the bird's size, overall shape, plurage and color-pattern. Include details on shape of bill, wings and tail. Give paticular emphasis to the marks you used to identify the bird. Reference to its resemblance in field guide descriptions is not enough. Describe that you actually saw in the field:

The size + shape ga Bullocks Orive, but somewhat plumps. Black head, lye, bill, backs, breast, wings, + tare; bright yellow belly, seides, under tare coverts + for along base of tare, rump, + part way down sides of tare. Fluis latter patiene was glumpsed only once, totongen the down sides of tare. Fluis latter patiene was glumpsed only once, totongen the DOX60's, because the bird stages so light + clicket spread its tare. There was also a bright yellow paties on shoulder, + a while way ber. Describe the bird's song or call, if heard, including method of delivery, i.e., from perch, in flight, etc: I haven't heard anything for years.

Describe the habitat in which you observed this bird: "Un'a big cottonwood at the near the South Platte River.

How were similarly appearing species eliminated? Black bladed is a dull greenen yellow & they color methodes the back. Male Baltimore is orang & has orange sides at the typ? tail. Scotts has bright yellow sides at base & tail. The black band tail. Scotts has bright yellow sides at base & tail. The black band

Prior experience with this and similarly appearing species: I have seen all of the Month american orioles & have seen several Scotts in Jerras + angona

Books, illustrations and advice consulted and how did these influence this documentation? Robbus, Brun & Z in in hand ishill il watched the bid

This report was written from notes made during after observation; from memory L PRINT NAME: <u>Thompsont</u> <u>G. Marsh</u> SIGNATURE: <u>High provide DATE</u>: 5-5-75 Street Address: <u>199 ash</u> Town and State: <u>Journ</u> <u>Cut</u> <u>ZIP: With</u> Return to: JACK REDEALL - <u>1450</u> South Alton Street, Englewood, Colorado 80110