23-89-19 A

Species: Little Gull, Larus minutus, juvenile plumage Date: 8/30/89, observed approximately 1:30 to 2:30 p.m. Location: Southwest corner of Cherry Creek Reservoir Other Observers: Another birdwatcher in the vicinity whom

I called over to see was Sam Palmer, 3327 Xanthis Court, Denver, Co. 80231.

Optical Equipment: 7 x 42 binoculars, Questar scope

- Distance: I viewed the bird while it was standing approximately 30 yards away.
- Field Marks: I first saw the bird as it was standing in amongst some Black Terns, compared to which it was bigger, and some Ring-Bill and California Gulls, compared to which it was significantly smaller. A very noticeable dark patch on top of the head was highlighted by a very white forehead. The dark patch on top of the head spread to the upper hind neck region. It had a dark eye with a dark spot that was approximately twice as large as the eye, which was separate from the eye on the side of its white face. The mantle was dark, and the darkness spread to the sides of the upper breast region, appearing as a partial breast band. The black cap was separated from the dark mantle by a white hind neck. On the dark-appearing sides of the uppers (lower scapular middle wing covert region) there were some faint pale vertical lines seen from approximately 30 yards away with the Questar. The wings were crossed and extended beyond the tail, and white could be seen on the outer part of the outer wing. When standing, a small, thin black tern-like bill was noticed, as well as pale reddish legs.

The bird flew frequently, and a dark leading edge to the upper outer wing was most noticeable. The dark upper inner wing coverts met on the lower back region. The trailing edge of the inner wing to the inner primaries did not have any dark trailing edge. This region appeared all white. The tail was white with a thin (approximately 25% of the tail) black terminal band. The underwings appeared all white.

Submitted by: Bob Righter, 2358 S. Fillmore, Denver, Co. 80210