CFO-ORC FILE # [997-5] 6-1 COLORADO FIELD ORNITHOLOGISTS - SIGHT RECORD FORM NEW STATE RECORD UNUSUAL RECORD Please print or type. Attach photographs and additional pages if necessary. Species: Lesser Nightlaute 1 chordeiles acutipen (Vernacular Name) (Scientific Name) 10 Sept. '95 Date(s): \_ Sex: 57 (?) Humber of birds seen: Plumage: adult ? Locality: e & Poudre R. Just s & Prospect ("Wren Site" Elevation: 5000 learest town: Fort Collins County: Lorimer Time(s) observed: 2 PM Duration of observation (e.g. 1 sec., 5 min., etc.): \_ 30 seconds to a minute Other observers who saw the bird with you (give names, addresses, telephone numbers): 4501 E Boardwork [970/225-1047] Ft collins 8:525 [970/225-1047] Dovid Ely Other observers who saw the bird independently (give names, addresses, telephone numbers) None that I know of Light conditions (position of bird to light and shade, direction and strength of light): Bird was above and then to the south & us, higher from the south Optical equipment used (e.g. power, types): 7×50 Nikor binocular Distance, and how measured: 100 07 If photographed, type of equipment and film: Who else photographed the bird? Describe the bird's behavior (e.g. flying, feeding, or habits used in identifying): Bird's odd herky-jerky flight pattern with much fluttering first cought our attention. Occasionally the bird would foll as if shot by a gun or "stalling", and then pull out of it. Flight direction in general was distinctly a to S as if migrating. Describe in detail the bird's relative size, overall shape, plumage and color pattern. Include details of bill, wing and tail shape. Give particular emphasis to marks you used to identify the bird. Reference to its resemblance to field guide descriptions and illustrations is not enough. Describe only what you actually observe in the field: The bird was obviously a night howk. It was a light buffy brown. Each wing was wanked with a bold white slash about midway between The "elbow" and the tip of the 1st primary. The wine tips were distinctly rounded making The wines overall appear shart compared to common nightbawks. The tail was notched and had a light band near, but not at the toil tip. First primary was shorter Than second. This helped provide the nonded splanged Look to the wing typ. Jack marks, flight pattern vos The wost distinctive (difficult To describe This adequately).

Description (continued):

Actorch unrelated, but perhops worthy of mention, kevin Cook & Forv collins claims to have seen a lesser nighthank at this same location in the late spring 91993 or 1994. (

Describe the bird's song and call, if given, including method of delivery, (i.e. from perch, in flight, duration):

Silent

Describe the general and specific habitat in which you observed the bird:

Ripanian comidor flanked by grovel ponds and flar terrain How did you eliminate similarly appearing species? Odd flight pattern, normded uniq tips & appearance of short wings, white wing sleshes slightly forther and toward mightips lighter coloration and broader tail band all were consistent with lesser vs Common. First primary was aborter than second - This is what made wing top "rowd" What is your prior experience with this and similarly appearing species? I have seen several lener Nightlawks in The Southwest US, one over The Cliffor Sever Ponds - 'S4 and one The min a Two Bruttes. I have seen thorsands of Common Nightlawks. This books, illustrations, recordings, other birders, etc. consulted and how this in-fluenced your identification:

a) at time of observation: None

when & Been in a line

b) after observation: Au the sta reference.

This report was written from reports made during  $\times$  , after observation  $\times$  from memory\_\_\_\_\_

Reporting observer: David Leatlenna Signature: David Leatlenna Address: 612 Stoven SY # 7 Ft collins, Co 80524 <u>970/416-0193(H)</u> 970/491-6303(w) All reports will be maintained permanently at the address below: Return to:

Bill Prather 13810 WCR 1 Longmont, CO 80504 303-776-2191