1999 - 133

CFO-RC FILE #_

COLORADO FIELD ORNITHOLOGISTS SIGHT RECORD FORM

Rcd 5/1/99 wpl

ACCEPT 7-0 RESULTS

PLEASE PRINT OR TYPE. ATTACH PHOTOGRAPHS AND ADDITIONAL PAGES IF NEC-ESSARY.

(COMMON NAME) (SCIENTIFIC NAME)

DATE(S) OF OCCURRENCE 5-2-99

NUMBER OF BIRDS SEEN _ ONE SEX MALE PLUMAGE AITERNATE

EXACT LOCATION NW were of LAMAR CENETREY

COUNTY Provides NEAREST TOWN LAMAR ELEVATION 3622

LENGTH OF OBSERVATION 5 mins TIME OF OBSERVATION ~7:00 am

OTHER OBSERVERS: BRANDON PORCIUMI, MARK JANOS (LITER) DAVE QUESENBEDRY VIL ZERBI Gene Rutherfued

LIGHT CONDITIONS - EARly Am son & clouds - crip, cool Light south wind

OPTICAL EQUIPMENT USED: ZIESS 10 x 40

DISTANCE FROM THE BIRD: 20 - 60 FEET

WAS THE BIRD PHOTOGRAPHED OF PHOTOGRAPHS SUBMITTED Y/N WHO TOOK THE PHOTOGRAPHS?

DESCRIBE THE GENERAL AND SPECIFIC HABITAT IN WHICH THE BIRD OCCURRED: CEMETARY with BIVE grass LAWN, MATURE PINE & EIM TREES No UNDER GROWTH DESCRIBE THE BIRD'S BEHAVIOR (FLYING; FEEDING; HABITS; BEHAVIOR USED IN IDENTIFYING THE SPECIES OR IN SUPPORTING THE ID):

BIRD WAS OBSERVED Foraging Amongst the PINE TREES, on the TRUNK, on The major Limbs and the outer Beanches

DESCRIBE THE BIRD IN DETAIL. THIS IS THE MOST IMPORTANT PART OF THIS RE-PORT. INCLUDE OVERALL SIZE AND SHAPE, AND HOW YOU PLACES THE BIRD IN THE GENERAL GROUP THAT YOU DID. GIVE COMPLETE DETAILS OF PLUMAGE, IN-CLUDING HEAD, THROAT, UNDERSIDES NAPE, BACK WINGS, AND TAIL. GIVE PAR-TICULAR EMPHASIS TO MARKS YOU USED TO ID. THE BIRD. DESCRIBE OTHER DE-TAILS SUCH AS COLOR AND SHAPE OF BILL, LEGS, EYES, ETC. THAT WERE IMPOR-TANT TO MAKING THE ID.. (ATTACH ADDITIONAL SHEETS, DRAWINGS, SKETCHES, OR COPY OF FIELD NOTES) :

- AVE SIZED WARBLER - that seem to tarage the Limbs HEAD - Bluish gray CROWN EXTENDING on to NECK. - Bill - BIALK - LONGER & PONTED - White SUPERCIllian EXTENDING to base of bill - EYE- BLACK. with Lower white CREST (1/2 ORBITAL Ring) - BLACK WIES EXTENDING DE KW ENE IN to ABRICULAR AREA ANd to SIDE of NECK · AURICULIT AREA BLACK WAS TERMINATE by A while PATCH on side of NECK. - YEILOW CHIN & THEORT BACK - UNMARKED Bluish- GRAY \$ Rump UNDER PERS . BREAST - BEILY - UNDER TAil CONERTS - White - Flanks - White with black STREAKing - A continuation of the black from the side of NECK. LEGS - WARK TAIL - DIO NOT PAY MUCH ATTIGATION - DARK

Wings - DARK with Two Wing BARS -, whitish outer EDGES of PRIMARIES DARELY NOTICABLE

a.

DISCUSS SIMILAR SPECIES AND HOW YOU ELIMINATED THEM:

GRAce's Weebler -> white check patch Audubais Warbler -> white check patch, black on BREAST & FLANKS No white supercillum

WHAT IS YOUR PRIOR EXPERIENCE WITH THIS AND SIMILARLY APPEARING

LIST BOOKS, GUIDES RECORDINGS, OR OTHER SOURCES CONSULTED AND HOW THIS INFLUENCED YOUR ID. (DURING AND AFTER OBSERVATION):

None consultan other than for scientific name

THIS REPORT WAS MADE FROMINOTES MADE DURING _____OR AFTER _____OBSERVATION, OR LATER FROM MEMORY_____.

DATE OF WRITTEN REPORT 5-3-99

REPORTING OBSERVER : RIC OLSON

SIGNATURE RO

9450 N. HEATHER DR. CASTLE ROCK, CO. 80104

303-790-7753

OLY@ECENTRAL.COM

YELLOW-THROATED WARBLER Dendroica dominica

1999-133 Led 1/2=/2 ipc ACCEPT 7-0

Location: Lamar Cemetery: Prowers County: Lamar, Colorado

Date: May 2-7, 1999

This bird was originally found by Brandon Percival.

Settina:

- -

This bird was seen by scores of Colorado birders, most notably much of a CFO field trip in early May. It was present, though hard to find, in the cemetery at Lamar. It was seen at various times, and under different conditions. I observed it on May 7th. The bird was fairly slow-moving and could be hard to find. It sometimes gleaned in the open and once we observed it perched on a phone wire (!). It barkgleaned, creeper-like, or moved out onto small limbs like a nuthatch. It stayed mainly in the upperparts of larger deciduous trees as well as large pines. I watched the bird at distances of 25-75 feet with B&L Elite 10x40 binoculars

Description:

It was a medium-small warbler, about 4-5" long. It was guite a striking warbler, patterned in white, black, gray, and yellow. It had a medium-sized to large bill, for a warbler, but it was long, thin and pointed. The bird's underparts were white. The flanks were extensively streaked with distinct black spots. The chin, throat and upper breast were a deep, clear yellow. The rest of the underparts were white. The upperparts were plain, dark gray. The wings were sooty gray with two large, distinct, white wingbars. The bird's face showed a complicated pattern, consisting of a black face patch, bordered by a white superciliary and lores above, and a white nape patch behind.

Similar Species:

The small size, slender pointed bill, white wingbars and active tree-top habits identify the bird as a warbler. Grace's is most similar but lacks the black face patch, the white nape spot and has yellow, not white, lores and superciliary. Blackburnian has yellow or orange face, streaked back and yellower undersides. Cape May lacks the black face patch, the white superciliary, and the clear yellow throat. Magnolia has a vellow belly and lacks the white nape spot. Townsend's lacks the white superciliary, the white nape spot and has black streaks on the breast.

Experience with Species:

I have seen all similar warblers with which this bird might be confused. I have seen Grace's Warbler in Colorado and elsewhere on several occasions, including in the spring of 1991, | have seen several previous Yellow-throated Warblers (three in Colorado), including several in Texas in March 1994

This report is prepared from notes made during and after observation.

Mark Janos 10 Sedum Ct. Pueblo, CO 81001

Martifanos