Scanned added

2000 - 9

DOCUMENTATION OF RARE BIRD SIGHTING (7-0 A) a led upl 5/20/00

Species: Little Blue Heron, Egretta caerulea

Date: 14 May 2000 Time seen: 4-4:30 pm

Locality: Havana Ponds, Rocky Mtn Arsenal, Adams Co.

Latilong: 39104 Optical equipment: Binocs, telescope 20-60 power

Number of birds seen: One Sex: Unknown Plumage: White immature

Light conditions: Good--cloudy bright

Distance from bird: 100-300 feet

Weather at time of observation. Bright cloudy, light wind off and on Prior weather & number of days since change: Generally good, 32 deg two nights before with a little rain

Habitat: reservoir edge

Behavior: First seen flying with a Snowy Egret to shoreline about 100 feet from our van. Moved, along edge of water, poking into water for food. Walk: plodding, deliberate, one-foot-at-a-time, not dainty, or rushed. During the observation, covered several hundred feet of shoreline before flying on to another section, where it continued the same behavior. First seen beside a Snowy Egret, later next to a Great Blue Heron.

Field marks observed: Heron by shape and behavior: long slightly curved neck, long legs, long bill.

 White plumage.

 Size:
 About the same size (height, not bulk) as the Snowy Egret beside it.

 Shape:
 Totally different shape from Snowy; LBH skinnier--slender heron with long crook neck that it held like a Great Blue yrather than a Snowy Egret, smooth feathering, no plumes or crest.

 Legs:
 Greenish gray.

 Bill:
 Bluish-gray bill with outer quarter dark. Bluish-gray lores extended up to eye, seemingly continuously from bill.

 Plumage:
 Overall white. Pale silvery-gray saddle. Shoulders smudged silvery-grayish. This gray tinge not conspicuous, and visibility depended on how bird stood with respect to the light.

Field marks not seen: This egret did not have dark edging on wings, except possibly one or two bars.

Similar species: Snowy Egret had plumes on back of head, held head and neck differently, had a chunkier shape and shorter neck, and different posture. Also yellow feet, black legs. Movement different from egrets. Bill and leg color distinguish from Snowy Egret & Reddish Egret. Size and bill color distinguish from white herons, etc.

Prior experience: Some, in Florida.

Description from: Notes taken in field

Observed by 10 members of Spring Count party.

ing & Aceph Jugery PO Box 584, Franktown CO 80

## 2000 - 9 \*